

Associations between the Wisconsin Inventory of Smoking Dependence Motives (WISDM-68) and Regional Brain Volumes in Adult Smokers

Alexander A. Brown BS^{1,2,3}, Roberto Cofresí PhD^{2,3}, Brett Froeliger PhD^{1,2,3}

¹Department of Psychiatry, University of Missouri, Columbia, MO, USA

²Department of Psychological Sciences, University of Missouri, Columbia, MO, USA

³Cognitive Neuroscience Systems Core Facility, University of Missouri, Columbia, MO, USA

Correspondence

Dr. Brett Froeliger

Departments of Psychiatry & Psychological Sciences

111 McAlester Hall

University of Missouri

Columbia, MO 65201

573-882-4785

froeligerb@health.missouri.edu

Accepted Manuscript

Abstract:

Introduction: The Wisconsin Inventory of Smoking Dependence Motives (WISDM-68) is a 68-item questionnaire to assess nicotine dependence as a multifactorial construct based on 13 theoretically derived smoking motives. Chronic smoking is associated with structural changes in brain regions implicated in the maintenance of smoking behavior; however, associations between brain morphometry and the various reinforcing components of smoking behavior remain unexamined. The present study investigated the potential association between smoking dependence motives and regional brain volumes in a cohort of 254 adult smokers.

Methods: The WISDM-68 was administered to participants at the baseline session. Structural magnetic resonance brain imaging (MRI) data from 254 adult smokers ($M_{\text{age}} = 42.7 \pm 11.4$) with moderate to severe nicotine dependence ($M_{\text{FTND}} = 5.4 \pm 2.0$) smoking for at least 2 years ($M_{\text{years}} = 24.3 \pm 11.8$) were collected and analyzed with Freesurfer.

Results: Vertex-wise cluster analysis revealed that high scores on the WISDM-68 composite, Secondary Dependence Motives (SDM) composite, and multiple SDM subscales were associated with lower cortical volume in the right lateral prefrontal cortex (cluster-wise p 's < 0.035). Analysis of subcortical volumes (i.e., nucleus accumbens, amygdala, caudate, pallidum) revealed several significant associations with WISDM-68 subscales, dependence severity (FTND), and overall exposure (pack years). No significant associations between cortical volume and other nicotine dependence measures or pack years were observed.

Conclusions: Results suggest that smoking motives may play a larger role in cortical abnormalities than addiction severity and smoking exposure *per se*, whereas subcortical volumes are associated with smoking motives, addiction severity, and smoking exposure.

Implications: The present study reports novel associations between the various reinforcing components of smoking behavior assessed by the WISDM-68 and regional brain volumes. Results suggest that the underlying emotional/cognitive/sensory processes that drive non-compulsive smoking behaviors may play a larger role in grey matter abnormalities of smokers than smoking exposure or addiction severity.

Accepted Manuscript

INTRODUCTION

Despite the well-known, detrimental effects of smoking tobacco, smoking continues to be the leading cause of preventable death and disease.¹ Why is smoking still so prevalent? The consensus is that nicotine dependence drives the maintenance of smoking behaviors and is largely responsible for smoking prevalence. Indeed, individuals with nicotine dependence are often unable to quit smoking, experience withdrawal symptoms after making a quit attempt, and continue to smoke despite awareness of the negative health consequences.

Neurobiologically, cigarette smokers with nicotine addiction have demonstrated abnormalities in brain structure such as decreased brain volume²⁻⁴ and accelerated cortical thinning^{5,6} compared to non-smokers.

Prevailing nicotine dependence assessments such as the Fagerström Test for Nicotine Dependence (FTND)⁷ and the Diagnostic and Statistical Manual of the American Psychiatric Association provide value by measuring nicotine dependence outcomes such as smoking heaviness and abstinence tolerability. FTND scores have shown to be predictive of smoking relapse among individuals making a quit attempt.⁸ However, smoking and nicotine reinforcement is a manifold process and most nicotine dependence assessments do not attempt to assess the underlying nature or mechanisms of nicotine dependence and do not investigate what characteristics of smoking may differentially reinforce the maintenance of smoking behavior.

In effort to elucidate the nature and behavioral mechanisms of nicotine addiction, Piper et al.⁹ developed the Wisconsin Inventory of Smoking Dependence Motives (WISDM-68), a 68-item questionnaire that attempts to define and measure dependence as a multifactorial construct on the basis of 13 theoretically derived smoking motives (see Supplementary Table A1). A follow-up study by Piper et al.¹⁰ performed latent class and factor analyses and

suggested that 4 subscales (Automaticity, Craving, Loss of Control, and Tolerance) represent the core features of dependence, and designated them as Primary Dependence Motives (PDM). The remaining 9 subscales (Affiliative Attachment, Behavioral Choice, Cognitive Enhancement, Cue Exposure/Associative Processes, Negative Reinforcement, Positive Reinforcement, Social/Environmental Goals, Taste/Sensory Properties, Weight Control) were designated as Secondary Dependence Motives (SDM) and are hypothesized to reflect the reasons people choose to smoke non-compulsively.

Multiple studies have found the PDM composite to be predictive of core nicotine dependence measures such as cigarettes per day (CPD), FTND scores, habitual or automatic motives for smoking cigarettes, and relapse.¹⁰⁻¹⁵ The SDM composite has shown to be a stronger predictor of instrumental and situational smoking and withdrawal-induced cravings and distress.^{10-12,14}

In an effort to further advance mechanistic understanding of the factors that reinforce the ongoing maintenance of smoking, the present study focused on examining how WISDM-68 smoking dependence motives are associated with brain morphometry. This is the first study to investigate neuroanatomical associations of smoking dependence motives.

Specifically, the primary goal was to examine and characterize the extent to which cortical and subcortical brain morphometry explained shared and unique variance in PDM and SDM. Provided the WISDM-68s concurrent validity with other dependence measures (FTND, CPD, pack years) that have shown to be associated with brain morphometry, we hypothesized that specific aspects smoking dependence motives would be differentially associated with brain volume. However, given the relative lack of extant literature in this domain, the pattern of associations were exploratory in nature. Using data from high resolution MRI brain scans, the present study utilized advanced surface-based morphometric analysis to assess the potential

relationship between WISDM-68 smoking dependence motives and regional brain volumes in a community sample of nicotine-dependent adult smokers.

METHODS

Participants

The data analyzed in the present study were compiled from seven smoking studies conducted at the University of Missouri-Columbia and the Medical University of South Carolina (MUSC). Primary findings of these studies are reported in prior publications.^{16–21} All studies were approved by the University of Missouri-Columbia and/or MUSC Internal Review Boards and were completed in accordance with the provisions of the World Medical Association Declaration of Helsinki. Informed consent was received from all participants prior to participation. Smokers ($n = 254$, $M_{\text{age}} = 42.7$ years ± 11.4) with moderate to severe nicotine addiction ($M_{\text{FTND}} = 5.4 \pm 2.0$) who had been smoking for at least 2 years ($M = 24.3$ years ± 11.8) were recruited from the local communities at both MUSC ($n = 151$) and the University of Missouri ($n = 103$).

The present study's cohort was gathered from previous smoking studies where various neuropsychiatric and substance use data were collected and utilized for inclusion/exclusion criteria. All subjects in the present study did not have a current, non-nicotinic substance use disorder. Subjects in the present study were administered the Center for Epidemiological Studies – Depression (CES-D), Beck's Anxiety Inventory (BAI), and were asked to report current or past psychosis, which if reported excluded participants from studies. 131 subjects (51.57%) of the present study's cohort were tested for current substance use and were administered the Mini-International Neuropsychiatric Interview (MINI). These 131 subjects did not test positive for current substance use and did not test positive for a psychiatric disorder when assessed by the MINI. The other 123 subjects in the present study's cohort were not excluded from their parent study if they reported or tested positive for current

substance use or reported a history of substance use disorder. Sample characteristics are detailed in Table 1.

Measures

The WISDM-68 and FTND were administered to all participants at their baseline session. Average CPD during the 30 days preceding the baseline session and number of years smoking were self-reported at the baseline session. Pack years was calculated as $(\text{CPD}/20) * (\text{years smoking})$. Scores for the 13 WISDM-68 subscales were calculated by taking the average score of all items belonging to the specified subscale. The WISDM-68 composite score was calculated as the sum of the 13 WISDM-68 subscales. The PDM composite score was calculated as the mean of Automaticity, Craving, Loss of Control, and Tolerance subscales. The SDM Composite was calculated as the mean score of Affiliative Attachment, Behavioral Choice/Amelioration, Cognitive Enhancement, Cue Exposure/Associative Processes, Negative Reinforcement, Positive Reinforcement, Social/Environmental Goals, Taste/Sensory Properties, and Weight Control subscales.

MRI data acquisition

3T MRI scanners (Siemens Prisma Fit – University of Missouri and MUSC [$n = 173$]; Siemens Tim Trio – MUSC [$n = 81$]), were used to acquire sets of high-resolution (1mm^3) T1-weighted structural brain images. Images were collected using standard T1-weighted magnetization prepared – rapid gradient echo (MPRAGE) pulse sequences (TR = 2300 ms [1900ms on Trio], TE = 2.26 ms, flip angle = 9° , 192 slices, 1mm^3 voxels, FOV = 256 mm). Participants were satiated and smoked a cigarette within 30 minutes of their MRI scan.

Data processing & analysis

Acquired T1s were visually inspected for quality assurance before being used as input for Freesurfer's (version 6.0.0) cortical reconstruction and volumetric segmentation pipelines. Participants with distorted T1s due to motion or artifact were excluded from the study. The

technical details of Freesurfer's cortical reconstruction and volumetric segmentation pipelines are described in prior publications. Once the cortical models were complete, a number of deformable procedures were performed for further data processing and analysis including surface inflation,²² registration to a spherical atlas which is based on individual cortical folding patterns to match cortical geometry across subjects,²³ parcellation of the cerebral cortex into units with respect to gyral and sulcal structure from the Desikan-Killiany (DK) atlas,²⁴ and creation of a variety of surface-based data including maps of curvature and sulcal depth. Critically, all Freesurfer-generated segmentations and cortical reconstructions were inspected and edited where needed to ensure accurate segmentations.

The potential relationship between cortical volume and nicotine dependence measures (WISDM-68, FTND, CPD) and smoking exposure (pack years) were examined using Freesurfer's surface-based, vertex-wise, general linear model cluster analyses, where general linear models are run at each vertex of the brain surface. Clusters in these analyses refer to brain regions where adjacent vertices share similar relationships at a specified statistical threshold for a specified general linear model.

For models assessing the relationship between cortical volume and nicotine dependence measures, each nicotine dependence measure was examined individually with age, education, sex, estimated total intracranial volumes (eTIV), and pack years entered as nuisance variables. Pack years was not included as a nuisance variable in models assessing the relationship between cortical volume and pack years. Cortical surface reconstructions for each participant were registered to Freesurfer's template brain – fsaverage – and were smoothed with a 10-millimeter full-width half-maximum (FWHM) gaussian spatial smoothing kernel. Clusters were corrected for multiple comparisons via permutation simulation ($n = 5000$) with a cluster forming threshold of $p < 0.05$. Only surviving clusters with a cluster-wise $p < 0.05$ were considered significant. Participant's total volumes from

significant resultant clusters were extracted and further analyzed using hierarchical linear regression models in SPSS (version 28.0). Partial correlations from the regression models are reported in the results.

The potential relationships between subcortical volumes, nicotine dependence measures, and smoking exposure were assessed using hierarchical linear regression models in SPSS. For models assessing the relationship between subcortical volumes and nicotine dependence measures, age, education, sex, eTIV, and pack years were entered as nuisance variables. Pack years was not included as a nuisance variable in models assessing the relationship between subcortical volumes and pack years. Partial correlations from the regression models are reported in the results.

RESULTS

Cortical volume analysis

Vertex-wise cluster analysis revealed significant negative associations between cortical volume and WISDM total, SDM composite, and SDM subscales (Affiliative Attachment, Behavioral Choice/Melioration, Cue Exposure/Associative Processes, Negative Reinforcement, Positive Reinforcement). Interestingly, all significant clusters were localized to the right lateral prefrontal cortex (IPFC) (Figure 1); specifically, the inferior frontal gyrus, rostral middle frontal, and lateral orbitofrontal cortex (all cluster-wise p 's < 0.035). The overlapping region of all clusters is displayed in Figure 1. Additional details of cluster analysis results are described in Table 2.

Each participant's total volume within each resultant cluster was extracted and used to further analyze the WISDM – cortical volume relationships via linear regression in SPSS (t [247] < -4.560, r < -0.278, p < 0.001 in all instances). Scatter plots displaying the residual volumes and residual WISDM scores of the significant associations are displayed in Supplementary

Figure A1. No significant associations were observed between cortical volume and PDM, FTND, CPD, or pack years.

Subcortical volume analysis

Primary Dependence Motives and subcortical volumes

High scores on the Automaticity subscale were associated with larger left nucleus accumbens volumes ($t [247] = 2.079, r = 0.131, p = 0.039$) and right nucleus accumbens volumes showed a similar trend ($t [247] = 1.868, r = 0.118, p = 0.063$). Tolerance showed a positive relationship with left caudate volumes ($t [247] = 2.05, r = 0.129, p = 0.041$).

Secondary Dependence Motives and subcortical volumes

Positive Reinforcement was negatively associated with right nucleus accumbens volumes ($t [247] = -2.179, r = -0.137, p = 0.03$). Taste/Sensory Properties showed both negative and positive associations with subcortical volumes. Bilateral nucleus accumbens and left amygdala volumes were negatively associated with Taste/Sensory Properties ($t [247] < -2.662, r < -0.166, p < 0.009$ in all instances), whereas bilateral pallidum volumes were positively associated with Taste/Sensory Properties ($t [247] > 2.910, r > 0.181, p < 0.005$ in both instances).

FTND, CPD, pack years, and subcortical volumes

FTND scores were positively associated with bilateral caudate volumes ($t [247] > 2.342, r > 0.146, p < 0.021$ in both instances). Pack Years was negatively associated with bilateral amygdala volumes ($t [248] < -2.273, r < -0.142, p < .025$ in both instances). No significant associations were observed between CPD and subcortical volumes. Significant subcortical volume associations are further detailed in Table 2.

DISCUSSION

The present study is the first to assess the relationship between WISDM-68 smoking dependence motives and brain morphometry. Primary findings include: 1) High scores on the WISDM-68 total, SDM composite, and multiple SDM subscales were associated with less cortical volume in the right lateral prefrontal cortex (IPFC); 2) Subcortical volumes (i.e., nucleus accumbens, amygdala, caudate, pallidum) revealed several significant associations with WISDM-68 subscales. Associations between subcortical volumes and PDM subscales were positive, whereas associations between subcortical volumes and SDM subscales were negative; 3) Amygdala volumes were negatively associated with pack years and caudate volumes were positively associated with FTND; 4) No significant associations between cortical volume and other nicotine dependence measures (FTND, CPD) or smoking exposure (pack years) were observed.

Cortical volume

Cluster analysis revealed significant negative associations between right lateral prefrontal cortex (IPFC) volumes and WISDM-68 total, SDM composite, and the SDM subscales Affiliative Attachment, Behavioral Choice/Melioration, Cue Exposure/Associative Processes, Negative Reinforcement, and Positive Reinforcement. Negative Reinforcement, a subscale representing the tendency or desire to smoke to ameliorate negative internal states, displayed the strongest association with cortical volume. The majority of the clusters were localized to the right inferior frontal gyrus (rIFG), a cortical region considered to be a locus of inhibitory control (IC)²⁵ and a prefrontal node of the hyperdirect pathway - a corticothalamic circuit involved in executing IC that has been shown to be dysregulated among individuals with nicotine dependence.^{16,17,26-28} Previous studies have shown smokers to exhibit less rIFG grey matter volume than non-smokers,^{4,29,30} and that smokers who relapsed following a quit attempt had less rIFG volume than those who remained abstinent.¹⁷ Additionally, a recent

study found that rIFG volume was positively associated with inhibitory control and rIFG thickness was positively associated with the ability to inhibit ad lib smoking during a smoking relapse analog task in a cohort of nicotine-dependent adult smokers.¹⁹

Interestingly, pack years, FTND, CPD, PDM composite, and PDM subscales were *not* significantly associated with cortical volumes. The lack of significant associations between cortical volume and FTND and CPD may be due in part to the homogeneity of the cohort as few participants reported low CPD and FTND scores. Additionally, many studies that have found associations between nicotine dependence severity and cortical volume have utilized ROI analyses which can be more liberal than surface-based, vertex-wise, whole brain analysis.

Although surprising, this is not the first imaging study of smokers to observe a lack of significant associations between FTND or pack years and brain volume. For example, Hanlon et al.³¹ found no associations between cortical volume and pack years, FTND, or years smoking when collapsing across age groups. Taken in conjunction with the volumetric associations observed in the present study, these results may suggest that alterations in cortical volume of smokers may reflect the progressive dysregulation of the cognitive/emotional processes that reinforce smoking behaviors. Future studies exploring the dissociable effects of smoking exposure, nicotine dependence, and smoking dependence motives on brain morphometry may provide valuable insight into the underlying mechanisms affecting brain morphometry in smokers.

Subcortical volumes

Automaticity

Nucleus accumbens volume was positively associated with the PDM subscale Automaticity. The nucleus accumbens is of particular interest in addiction research as the reinforcing effects of most drugs of abuse such as nicotine depend on dopamine release in the nucleus accumbens.³² The early stages of nicotine addiction are thought to be initiated by a nicotine-induced increase of dopamine release in the nucleus accumbens, but chronic nicotine exposure leads to glutamatergic mediated neuroplasticity in prefrontal - nucleus accumbens (i.e. corticostriatal) circuitry that in turn mediates cue-induced drug seeking.³³ As addiction becomes more entrenched, the glutamatergic corticostriatal projections become further dysregulated, and morphological changes emerge, such as increased dendritic spine density, synapse size, and dendritic branching.^{32,34} Given that the Automaticity subscale captures smoking without conscious intention or awareness, a behavioral pattern representative of "end-stage addiction,"³³ and was significantly correlated with pack years, FTND, and CPD, the positive association between Automaticity and nucleus accumbens volume may be attributed to changes in glutamatergic-mediated synaptic plasticity and dendritic arborization in corticostriatal circuitry. Indeed, findings from non-addiction studies suggest that changes in structural complexity via dendritic arborization and synaptic plasticity, independent of changes in neuronal count, may contribute to changes in gray matter volume.³⁵⁻³⁷ However, in contrast, see.³⁸ More research is needed to better understand how nucleus accumbens volume is associated with automatic or compulsive smoking behavior and cessation outcomes following a quit attempt.³⁹

Positive Reinforcement

In contrast to the positive association between the Automaticity subscale and nucleus accumbens volume, the Positive Reinforcement subscale was *negatively* associated with nucleus accumbens volume. The Positive Reinforcement subscale captures the motivation to experience a positive consequence of smoking, starkly different than the Automaticity subscale, which captures smoking compulsively and automatically without consideration of consequences. Indeed, Automaticity scores were positively correlated with CPD and pack years, whereas Positive Reinforcement scores were not. As addiction progresses, the subjective experience of "rewarding" positive affective states from smoking is confounded with the reversal from a nicotine withdrawal state. Thus, the desire to experience a positive consequence of smoking may be a motive among smokers who have yet to enter end-stage addiction, or among highly dependent smokers, the experience of relief from withdrawal. Future longitudinal studies are needed that conduct a principled examination of a broad range of nicotine and tobacco exposure measures: dose, chronicity, and duration to further disentangle these associations.

FTND and Tolerance

FTND and Tolerance were positively associated with caudate volumes. Of all WISDM composites and subscales, Tolerance showed the strongest correlation with FTND ($r = 0.610$), and was significantly correlated with pack years and CPD. The caudate contains a high density of dopamine receptors and prior research has observed increased dopamine uptake in the caudate of smokers compared to non-smokers.⁴⁰ The caudate plays a functional role in executive function and shows significant functional connectivity with other brain regions implicated in smoking such as the anterior cingulate, insula, thalamus, and inferior frontal and middle frontal gyri.⁴¹ Indeed, an fMRI study by Feng et al.⁴² found weaker resting state functional connectivity between the caudate and anterior cingulate in smokers compared

to non-smokers, which was associated with cognitive control. Previous findings regarding smoking and caudate morphometry are limited. However, a study by Li et al.⁴³ observed larger caudate volumes in smokers compared to non-smokers. Given that the Tolerance subscale captures the need to smoke increasing amounts over time and Tolerance scores were strongly correlated with FTND, CPD, and pack years, the positive associations between caudate volume and FTND and Tolerance may be driven by the previously described changes to corticostriatal projections associated with end-stage addiction.

Pack Years

The present study's finding of a negative association between amygdala volumes and pack years are similar to the findings reported in Durazzo et al.³⁸ that observed a negative association between pack years and amygdala volume in adult smokers. Previous studies have also found smokers to exhibit smaller amygdala volumes in comparison to non-smokers.³¹ The amygdala contains a high density of nicotinic acetylcholine receptors (nAChR)⁴⁴ and a previous rodent study by Huang et al.⁴⁵ found that chronic nicotine exposure promoted long-lasting synaptic changes in the amygdala. However, it remains unclear if nicotine-associated changes in cholinergic receptors or synapses are related to amygdala morphometry in smokers.

Taste/Sensory Properties

Bilateral nucleus accumbens and left amygdala volumes were negatively associated with Taste/Sensory Properties, whereas bilateral pallidum volumes showed a positive association with Taste/Sensory Properties. The pallidum is a relatively dopamine-rich structure with moderate expression of dopamine receptors and receives dense, efferent γ -aminobutyric acid (GABA) and peptide projections from the nucleus accumbens that are critical for expressing motivation behaviors.⁴⁶ A recent study by Zou et al.⁴⁷ found that 10 year-old children whose

fathers smoked periconceptually had larger pallidum volumes than age-matched children whose parents did not smoke.

The nucleus accumbens, amygdala, and pallidum are structures implicated in gustatory conditioning, associative learning, and taste aversion (for review see ⁴⁸). Although nicotine is the primary reinforcer of smoking addiction it is not the only reinforcer, as indicated by previous studies showing the low success rates of nicotine replacement therapy and the minimal enjoyment smokers report when dosed with nicotine infusions.⁴⁹ Therefore, non-nicotine components of smoking such as sensory properties (i.e., taste, smell, respiratory tract stimulation) play a significant role in smoking reinforcement. For example, local anesthesia of the airways significantly reduces smoking satisfaction,⁵⁰ and attenuation of olfactory/taste cues diminishes enjoyment and behavioral reinforcement effects of cigarette smoke.⁵¹ More research is needed to better understand the neurobiological mechanisms of how taste and sensory properties of smoking are associated with brain morphometry.

Limitations and Future Directions

Provided the lack of genetic, environmental, and longitudinal data for this study, it cannot be conclusively determined if the observed associations between smoking and brain morphometry are caused by smoking-related effects or are due to pre-existing genetic/environmental factors that alter brain morphometry. Indeed, previous research has shown that cortical volume is associated with certain genetic polymorphisms.^{53,54} In the case that smoking did not impact brain morphometry, one may interpret the results as potential pre-disposed risk factors for developing nicotine dependence (e.g., individuals with less right IPFC volume are more likely to develop nicotine dependence). Longitudinal brain imaging studies of smokers are needed to better understand the effects of smoking and nicotine exposure on brain morphometry.

The present study's data were compiled from multiple studies across different sites, and included slight variation in scanning protocols. However, all T1s were visually inspected before processing and no observable differences were apparent. Distorted T1s due to motion or artifact were excluded from the study. Additionally, all Freesurfer-generated segmentations and cortical reconstructions were inspected and edited where needed to ensure accurate segmentations. Questionnaires were administered the same across studies, but variation due to different lab environments and personnel across sites may have affected the results.

The present study presents a novel, retrospective analysis of associations between the WISDM-68 and brain morphometry. Future prospective studies assessing the relationship between brain morphometry and the WISDM-68 may consider alternative statistical approaches than the ones used here. For example, Piper et al.'s study¹⁰ describing the latent class and factor analyses for the formulation of PDM and SDM found that after controlling for the variance of PDM, SDM was less or non-predictive of many tobacco dependence measures whereas controlling for the variance of SDM when assessing PDM's predictive value had minimal effect. However, given the novelty of the present study and the lack of clarity regarding the relationship between nicotine dependence and brain morphometry, we chose to only include covariates in our models that have been shown to influence brain morphometry (age, education, sex, pack years).

The present study's cohort had a wide age range of participants (20-66 years old), but most were middle-aged adults with extensive smoking histories. A younger cohort of recently dependent smokers may yield better insight into how smoking dependence motives are related to brain morphometry and the development of nicotine addiction.

CONCLUSIONS

The present study presents novel findings on associations between regional brain volumes and WISDM-68 smoking dependence motives. Significant associations between cortical volumes and WISDM-68 total, SDM composite, and SDM subscales were localized to the right IPFC. Cortical volumes were not significantly associated with WISDM PDM composite or subscales, FTND, CPD, or pack years. Analysis of subcortical volumes revealed several significant associations with WISDM-68 subscales. Associations between subcortical volumes and WISDM PDM subscales were positive, whereas associations between subcortical volumes and WISDM SDM subscales were negative. Together, these results suggest that grey matter abnormalities play a significant role in the emotional/cognitive/sensory processes that reinforce smoking, as opposed to simply anatomical correlates of general exposure to smoking or addiction severity. Longitudinal brain imaging studies of smokers are needed to better understand the relationship between brain morphometry and smoking exposure, addiction severity, and smoking dependence motives.

Accepted Manuscript

ACKNOWLEDGEMENTS:

This work was supported by research grants from the National Institute of Health (NIH) / National Institute on Drug Abuse (NIDA): R01DA048094 [BF], R01DA038700 [BF], R01DA033459 [BF], UG3DA048510 [BF], P30CA138313 [MUSC Hollings Cancer Center].

AUTHORS CONTRIBUTION

BF was responsible for the study concept and design. AAB performed the data analysis and drafted the manuscript. BF and RC provided critical revision of the manuscript for important intellectual content. All authors critically reviewed the content and approved the final version of the manuscript submitted for publication.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study may be available from the corresponding author (BF), upon reasonable request. The data are not publicly available due to restrictions related to internal review board policies and informed consent limitations under which the data were originally collected.

CONFLICT OF INTEREST

AAB, RC, and BF reported no biomedical financial interests or potential conflicts of interest.

ETHICS APPROVAL AND PATIENT CONSENT STATEMENT

The present study was approved by the University of Missouri and Medical University of South Carolina Internal Review Boards and was carried out in accordance with the provisions of the World Medical Association Declaration of Helsinki. Informed consent was obtained for all individuals prior to participation.

References

1. CDC. Current Cigarette Smoking Among Adults in the United States. Centers for Disease Control and Prevention. Published March 16, 2022. Accessed August 17, 2022. https://www.cdc.gov/tobacco/data_statistics/fact_sheets/adult_data/cig_smoking/index.htm
2. Durazzo TC, Cardenas VA, Studholme C, Weiner MW, Meyerhoff DJ. Non-treatment-seeking heavy drinkers: Effects of chronic cigarette smoking on brain structure. *Drug Alcohol Depend.* 2007;87(1):76-82. doi:10.1016/j.drugalcdep.2006.08.003
3. Sutherland MT, Riedel MC, Flannery JS, et al. Chronic cigarette smoking is linked with structural alterations in brain regions showing acute nicotinic drug-induced functional modulations. *Behav Brain Funct.* 2016;12(1):16. doi:10.1186/s12993-016-0100-5
4. Gallinat J, Meisenzahl E, Jacobsen LK, et al. Smoking and structural brain deficits: a volumetric MR investigation. *Eur J Neurosci.* 2006;24(6):1744-1750. doi:10.1111/j.1460-9568.2006.05050.x
5. Durazzo TC, Meyerhoff DJ, Yoder KK. Cigarette smoking is associated with cortical thinning in anterior frontal regions, insula and regions showing atrophy in early Alzheimer's Disease. *Drug Alcohol Depend.* 2018;192:277-284. doi:10.1016/j.drugalcdep.2018.08.009
6. Karama S, Ducharme S, Corley J, et al. Cigarette smoking and thinning of the brain's cortex. *Mol Psychiatry.* 2015;20(6):778-785. doi:10.1038/mp.2014.187
7. Heatherton TF, Kozlowski LT, Frecker RC, Fagerstrom KO. The Fagerström Test for Nicotine Dependence: a revision of the Fagerstrom Tolerance Questionnaire. *Br J Addict.* 1991;86(9):1119-1127. doi:10.1111/j.1360-0443.1991.tb01879.x
8. Fagerström K, Russ C, Yu CR, Yunis C, Foulds J. The Fagerström Test for Nicotine Dependence as a predictor of smoking abstinence: a pooled analysis of varenicline clinical trial data. *Nicotine Tob Res Off J Soc Res Nicotine Tob.* 2012;14(12):1467-1473. doi:10.1093/ntr/nts018
9. Piper ME, Piasecki TM, Federman EB, et al. A Multiple Motives Approach to Tobacco Dependence: The Wisconsin Inventory of Smoking Dependence Motives (WISDM-68). *J Consult Clin Psychol.* 2004;72(2):139-154. doi:10.1037/0022-006X.72.2.139
10. Piper ME, Bolt DM, Kim SY, et al. Refining the tobacco dependence phenotype using the Wisconsin Inventory of Smoking Dependence Motives (WISDM). *J Abnorm Psychol.* 2008;117(4):747-761. doi:10.1037/a0013298
11. Baker TB, Piper ME, Schlam TR, et al. Are tobacco dependence and withdrawal related amongst heavy smokers? Relevance to conceptualizations of dependence. *J Abnorm Psychol.* 2012;121:909-921. doi:10.1037/a0027889
12. Piasecki TM, Piper ME, Baker TB. Refining the tobacco dependence phenotype using the Wisconsin Inventory of Smoking Dependence Motives: II. Evidence from a laboratory self-administration assay. *J Abnorm Psychol.* 2010;119:513-523. doi:10.1037/a0020235

13. Shenassa ED, Graham AL, Burdzovic JA, Buka SL. Psychometric properties of the Wisconsin Inventory of Smoking Dependence Motives (WISDM-68): A replication and extension. *Nicotine Tob Res.* 2009;11(8):1002-1010. doi:10.1093/ntr/ntp109
14. Piasecki TM, Piper ME, Baker TB, Hunt-Carter EE. WISDM primary and secondary dependence motives: Associations with self-monitored motives for smoking in two college samples. *Drug Alcohol Depend.* 2011;114(2):207-216. doi:10.1016/j.drugalcdep.2010.10.005
15. Parrott CE, Rathnayaka N, Blalock JA, et al. Examination of the Wisconsin Inventory of Smoking Dependence Motives (WISDM-68) Factor Structure in a Sample of Pregnant Smokers. *Nicotine Tob Res.* 2015;17(6):653-660. doi:10.1093/ntr/ntu238
16. Bell S, Froeliger B. Associations Between Smoking Abstinence, Inhibitory Control, and Smoking Behavior: An fMRI Study. *Front Psychiatry.* 2021;12. Accessed August 1, 2022. <https://www.frontiersin.org/articles/10.3389/fpsy.2021.592443>
17. Froeliger B, McConnell PA, Bell S, et al. Association Between Baseline Corticothalamic-Mediated Inhibitory Control and Smoking Relapse Vulnerability. *JAMA Psychiatry.* 2017;74(4):379-386. doi:10.1001/jamapsychiatry.2017.0017
18. Froeliger B, Mathew AR, McConnell PA, et al. Restructuring Reward Mechanisms in Nicotine Addiction: A Pilot fMRI Study of Mindfulness-Oriented Recovery Enhancement for Cigarette Smokers. *Evid-Based Complement Altern Med ECAM.* 2017;2017:7018014. doi:10.1155/2017/7018014
19. Brown AA, Upton S, Craig S, Froeliger B. Associations between right inferior frontal gyrus morphometry and inhibitory control in individuals with nicotine dependence. *Drug Alcohol Depend.* 2023;244:109766. doi:10.1016/j.drugalcdep.2023.109766
20. Newman-Norlund RD, Gibson M, McConnell PA, Froeliger B. Dissociable Effects of Theta-Burst Repeated Transcranial Magnetic Stimulation to the Inferior Frontal Gyrus on Inhibitory Control in Nicotine Addiction. *Front Psychiatry.* 2020;11. Accessed August 1, 2022. <https://www.frontiersin.org/articles/10.3389/fpsy.2020.00260>
21. Froeliger B, McConnell PA, Stankeviciute N, McClure EA, Kalivas PW, Gray KM. The effects of N-Acetylcysteine on frontostriatal resting-state functional connectivity, withdrawal symptoms and smoking abstinence: A double-blind, placebo-controlled fMRI pilot study. *Drug Alcohol Depend.* 2015;156:234-242. doi:10.1016/j.drugalcdep.2015.09.021
22. Fischl B, Sereno MI, Dale AM. Cortical Surface-Based Analysis: II: Inflation, Flattening, and a Surface-Based Coordinate System. *NeuroImage.* 1999;9(2):195-207. doi:10.1006/nimg.1998.0396
23. Fischl B, Sereno MI, Tootell RBH, Dale AM. High-resolution intersubject averaging and a coordinate system for the cortical surface. *Hum Brain Mapp.* 1999;8(4):272-284. doi:10.1002/(SICI)1097-0193(1999)8:4<272::AID-HBM10>3.0.CO;2-4
24. Desikan RS, Ségonne F, Fischl B, et al. An automated labeling system for subdividing the human cerebral cortex on MRI scans into gyral based regions of interest. *NeuroImage.* 2006;31(3):968-980. doi:10.1016/j.neuroimage.2006.01.021

25. Aron AR, Robbins TW, Poldrack RA. Inhibition and the right inferior frontal cortex: one decade on. *Trends Cogn Sci*. 2014;18(4):177-185. doi:10.1016/j.tics.2013.12.003
26. Froeliger B, Modlin LA, Kozink RV, et al. Frontoparietal Attentional Network Activation Differs Between Smokers and Nonsmokers during Affective Cognition. *Psychiatry Res*. 2013;211(1):57-63. doi:10.1016/j.psychres.2012.05.002
27. Froeliger B, Modlin L, Wang L, Kozink RV, McClernon FJ. Nicotine withdrawal modulates frontal brain function during an affective Stroop task. *Psychopharmacology (Berl)*. 2012;220(4):707-718. doi:10.1007/s00213-011-2522-y
28. Kozink RV, Lutz AM, Rose JE, Froeliger B, McClernon FJ. Smoking withdrawal shifts the spatiotemporal dynamics of neurocognition. *Addict Biol*. 2010;15(4):480-490. doi:10.1111/j.1369-1600.2010.00252.x
29. Brody AL, Mandelkern MA, Jarvik ME, et al. Differences between smokers and nonsmokers in regional gray matter volumes and densities. *Biol Psychiatry*. 2004;55(1):77-84. doi:10.1016/s0006-3223(03)00610-3
30. Fritz HC, Wittfeld K, Schmidt CO, et al. Current Smoking and Reduced Gray Matter Volume— a Voxel-Based Morphometry Study. *Neuropsychopharmacology*. 2014;39(11):2594-2600. doi:10.1038/npp.2014.112
31. Hanlon CA, Owens MM, Joseph JE, et al. Lower subcortical gray matter volume in both younger smokers and established smokers relative to non-smokers. *Addict Biol*. 2016;21(1):185-195. doi:10.1111/adb.12171
32. Volkow ND, Michaelides M, Baler R. The Neuroscience of Drug Reward and Addiction. *Physiol Rev*. 2019;99(4):2115-2140. doi:10.1152/physrev.00014.2018
33. Kalivas PW, Volkow ND. The Neural Basis of Addiction: A Pathology of Motivation and Choice. *Am J Psychiatry*. 2005;162(8):1403-1413. doi:10.1176/appi.ajp.162.8.1403
34. Robinson TE, Gorny G, Mitton E, Kolb B. Cocaine self-administration alters the morphology of dendrites and dendritic spines in the nucleus accumbens and neocortex. *Synapse*. 2001;39(3):257-266. doi:10.1002/1098-2396(20010301)39:3<257::AID-SYN1007>3.0.CO;2-1
35. Freeman SH, Kandel R, Cruz L, et al. Preservation of Neuronal Number Despite Age-Related Cortical Brain Atrophy in Elderly Subjects Without Alzheimer Disease. *J Neuropathol Exp Neurol*. 2008;67(12):1205-1212. doi:10.1097/NEN.0b013e31818fc72f
36. Lerch JP, Yiu AP, Martinez-Canabal A, et al. Maze training in mice induces MRI-detectable brain shape changes specific to the type of learning. *NeuroImage*. 2011;54(3):2086-2095. doi:10.1016/j.neuroimage.2010.09.086
37. Scheibel ME, Lindsay RD, Tomiyasu U, Scheibel AB. Progressive dendritic changes in aging human cortex. *Exp Neurol*. 1975;47(3):392-403. doi:10.1016/0014-4886(75)90072-2

38. Durazzo TC, Meyerhoff DJ, Yoder KK, Murray DE. Cigarette smoking is associated with amplified age-related volume loss in subcortical brain regions. *Drug Alcohol Depend.* 2017;177:228-236. doi:10.1016/j.drugalcdep.2017.04.012
39. Froeliger B, Kozink RV, Rose JE, Behm FM, Salley AN, McClernon FJ. Hippocampal and striatal gray matter volume are associated with a smoking cessation treatment outcome: results of an exploratory voxel-based morphometric analysis. *Psychopharmacology (Berl).* 2010;210(4):577-583. doi:10.1007/s00213-010-1862-3
40. Salokangas RKR, Vilkinen H, Ilonen T, et al. High Levels of Dopamine Activity in the Basal Ganglia of Cigarette Smokers. *Am J Psychiatry.* 2000;157(4):632-634. doi:10.1176/appi.ajp.157.4.632
41. Robinson JL, Laird AR, Glahn DC, et al. The functional connectivity of the human caudate: An application of meta-analytic connectivity modeling with behavioral filtering. *NeuroImage.* 2012;60(1):117-129. doi:10.1016/j.neuroimage.2011.12.010
42. Feng D, Yuan K, Li Y, et al. Intra-regional and inter-regional abnormalities and cognitive control deficits in young adult smokers. *Brain Imaging Behav.* 2016;10(2):506-516. doi:10.1007/s11682-015-9427-z
43. Li Y, Yuan K, Cai C, et al. Reduced frontal cortical thickness and increased caudate volume within fronto-striatal circuits in young adult smokers. *Drug Alcohol Depend.* 2015;151:211-219. doi:10.1016/j.drugalcdep.2015.03.023
44. Mineur YS, Fote GM, Blakeman S, Cahuzac ELM, Newbold SA, Picciotto MR. Multiple Nicotinic Acetylcholine Receptor Subtypes in the Mouse Amygdala Regulate Affective Behaviors and Response to Social Stress. *Neuropsychopharmacology.* 2016;41(6):1579-1587. doi:10.1038/npp.2015.316
45. Huang YY, Kandel ER, Levine A. Chronic nicotine exposure induces a long-lasting and pathway-specific facilitation of LTP in the amygdala. *Learn Mem.* 2008;15(8):603-610. doi:10.1101/lm.975308
46. Mogenson GJ, Brudzynski SM, Wu M, Yang CR, Yim CCY. From Motivation to Action: A Review of Dopaminergic Regulation of Limbic → Nucleus Accumbens → Ventral Pallidum → Pedunculopontine Nucleus Circuitries Involved in Limbic-Motor Integration. In: *Limbic Motor Circuits and Neuropsychiatry.* CRC Press; 1993.
47. Zou R, Boer OD, Felix JF, et al. Association of Maternal Tobacco Use During Pregnancy With Preadolescent Brain Morphology Among Offspring. *JAMA Netw Open.* 2022;5(8):e2224701. doi:10.1001/jamanetworkopen.2022.24701
48. Yamamoto T, Ueji K. Brain Mechanisms of Flavor Learning. *Front Syst Neurosci.* 2011;5. Accessed October 6, 2022. <https://www.frontiersin.org/articles/10.3389/fnsys.2011.00076>
49. Rose JE, Behm FM, Westman EC, Bates JE, Salley A. Pharmacologic and sensorimotor components of satiation in cigarette smoking. *Pharmacol Biochem Behav.* 2003;76(2):243-250. doi:10.1016/j.pbb.2003.07.002

50. Rose JE, Tashkin DP, Ertle A, Zinser MC, Lafer R. Sensory blockade of smoking satisfaction. *Pharmacol Biochem Behav.* 1985;23(2):289-293. doi:10.1016/0091-3057(85)90572-6
51. Perkins KA, Gerlach D, Vender J, Meeker J, Hutchison S, Grobe J. Sex differences in the subjective and reinforcing effects of visual and olfactory cigarette smoke stimuli. *Nicotine Tob Res.* 2001;3(2):141-150. doi:10.1080/14622200110043059
52. Smith SS, Piper ME, Bolt DM, et al. Development of the Brief Wisconsin Inventory of Smoking Dependence Motives. *Nicotine Tob Res.* 2010;12(5):489-499. doi:10.1093/ntr/ntq032
53. Mon A, Durazzo TC, Gazdzinski S, Hutchison KE, Pennington D, Meyerhoff DJ. Brain-derived neurotrophic factor genotype is associated with brain gray and white matter tissue volumes recovery in abstinent alcohol-dependent individuals. *Genes Brain Behav.* 2013;12(1):98-107. doi:10.1111/j.1601-183X.2012.00854.x
54. Pezawas L, Verchinski BA, Mattay VS, et al. The Brain-Derived Neurotrophic Factor val66met Polymorphism and Variation in Human Cortical Morphology. *J Neurosci.* 2004;24(45):10099-10102. doi:10.1523/JNEUROSCI.2680-04.2004

Accepted Manuscript

Table 1. Sample Characteristics**Demographics**

Sample Size (Female)	254 (126)
Race - N (%)	
White	185 (72.8%)
Black	57 (22.4%)
Other	12 (4.7%)
Age - mean (SD)	42.7 (11.4)
Education - mean (SD)	14.1 (2.2)

Baseline Measures - mean (SD)

Nicotine Dependence (FTND score)	5.4 (2.0)
Pack Years	22.5 (16.0)
Years Smoking	24.3 (11.8)
Cigarettes Per Day (30-day average)	17.6 (7.5)

WISDM Composites and Subscale Scores - mean (SD)

Total ^a	56.0 (15.4)
Primary Dependence Motives ^b	4.4 (1.3)
Secondary Dependence Motives ^c	3.8 (1.1)
Affiliative Attachment	2.9 (1.6)
Automaticity	4.0 (1.7)
Loss of Control	4.4 (1.6)
Behavioral Choice/Melioration	3.4 (1.4)
Cognitive Enhancement	3.6 (1.6)
Craving	4.5 (1.5)
Cue Exposure/Associative Processes	4.6 (1.3)
Negative Reinforcement	4.5 (1.5)
Positive Reinforcement	4.1 (1.6)
Social/Environmental Goals	3.7 (1.9)
Taste/Sensory Properties	4.3 (1.5)
Tolerance	4.6 (1.4)
Weight Control	2.8 (1.6)

^aWISDM Total Score = Sum of the 13 WISDM subscales

^bPrimary Dependence Motives = Mean of Automaticity, Loss of Control, Craving, and Tolerance

^cSecondary Dependence Motives = Mean of Affiliative Attachment, Behavioral Choice/Melioration, Cognitive Enhancement, Cue Exposure/Associative Processes, Negative Reinforcement, Positive Reinforcement, Social/Environmental Goals, Taste, and Weight Control

Accepted Manuscript

Table 2. Cluster Analysis Results

WISDM Scale	Peak Region	MNI Coordinates of Peak Vertex			Peak Vertex <i>t</i> -Statistic	Cluster-Wise <i>p</i>	Cluster-Wise <i>p</i> 90% Confidence Interval	Cluster Size (mm ²)
		X	Y	Z				
Total	Pars Orbitalis	46	36.9	-9.6	-4.941	0.025	0.023 - 0.028	2055.80
Secondary Dependence Motives	Pars Orbitalis	46	36.9	-9.6	-4.907	0.014	0.012 - 0.016	2482.07
Affiliative Attachment	Pars Orbitalis	46.9	35.5	-9.2	-3.698	0.033	0.030 - 0.037	1894.20
Behavioral Choice/Melioration	Pars Orbitalis	46	36.9	-9.6	-4.379	0.034	0.031 - 0.037	1901.59
Cue Exposure/Associative Processes	Pars Orbitalis	46.4	35.6	-8.2	-5.29	0.033	0.030 - 0.037	1878.58
Negative Reinforcement	Pars Orbitalis	46	36.9	-9.6	-6.177	0.007	0.006 - 0.009	2840.88
Positive Reinforcement	Pars Orbitalis	46.5	37.7	-10.9	-6.609	0.022	0.019 - 0.024	2164.31

Table 3. Subcortical Volume Associations

Subcortical Structure	Associated Measure	Direction	t	Partial Correlation	p
Right Nucleus Accumbens	Positive Reinforcement	Negative	-2.179	-0.137	0.03
	Taste/Sensory Properties	Negative	-3.266	-0.203	0.001
Left Nucleus Accumbens	Taste/Sensory Properties	Negative	-3.946	-0.243	< 0.001
	Automaticity	Positive	2.079	0.131	0.039
Right Amygdala	Pack Years	Negative	-2.274	-0.143	0.024
Left Amygdala	Pack Years	Negative	-3.126	-0.195	0.002
	Taste/Sensory Properties	Negative	-2.663	-0.167	0.008
Right Caudate	FTND	Positive	2.531	0.159	0.012
Left Caudate	FTND	Positive	2.343	0.147	0.02
	Tolerance	Positive	2.05	0.129	0.041
Right Pallidum	Taste/Sensory Properties	Positive	3.091	0.193	0.002
Left Pallidum	Taste/Sensory Properties	Positive	2.911	0.182	0.004

Figure 1. Associations Between Cortical Volume and WISDM-68




Figure 1. Corrected cluster analysis results (blue) overlaid on the Freesurfer template (fsaverage) inflated surface displaying the negative associations between cortical volume and WISDM composite and subscale scores controlling for age, education, sex, estimated total intracranial volume, and pack years (all cluster-wise p -values < 0.035). The overlapping region of all clusters is displayed on the bottom right. Outlines of the right inferior frontal gyrus (black), rostral middle frontal (yellow), and lateral orbitofrontal cortex (white) from the Desikan-Killiany atlas are also overlaid on the inflated surface.